

Dynamical Systems, Topology and Conductivity in Normal Metals in strong magnetic fields.

A.Ya.Maltsev¹, S.P.Novikov^{1,2}.

⁽¹⁾ L.D.Landau Institute for Theoretical Physics,
119334 ul. Kosygina 2, Moscow,
maltsev@itp.ac.ru , novikov@itp.ac.ru

⁽²⁾ IPST, University of Maryland,
College Park MD 20742-2431,USA
novikov@ipst.umd.edu

Abstract

We represent here the full description of all asymptotic regimes of conductivity behavior in the so-called "Geometric Strong Magnetic Field limit" in the 3D single crystal normal metals with topologically complicated Fermi surfaces. In particular, new observable integer-valued characteristics of conductivity of the topological origin were introduced by the present authors few years ago; they are based on the Topological Resonance found by the present authors and play the basic role in the total picture. Our investigation is based on the study of dynamical systems on Fermi surfaces for the semi-classical motion of electron in magnetic field realized by the Moscow topological group.

This paper is the reduced and improved version of the paper cond-mat/0304471. The authors are very grateful to Prof. Joel L. Lebowitz for his help and interest to this work.

1 Introduction.

We are going to consider the implications of the so called "Geometric Strong Magnetic Field limit" for the conductivity in normal metals with topologically complicated Fermi surface in the presence of the homogeneous magnetic field. The corresponding limit can be defined by the relation $\omega_B \tau \gg 1$. Here ω_B is the cyclotron frequency for the electron in crystal and τ is the mean free motion time between the scattering acts. This means actually that this theory is based on the Kinetic equation for the electron gas in crystal for the semiclassical electrons in the external fields. Let us say that the corresponding conditions for the external fields are always satisfied for the experimentally available electric and magnetic fields in the case of normal metals. We can speak, for example, about the limit of very strong magnetic fields in the experimental sense where the semiclassical approximation still gives the main features of transport phenomena. It works until the magnetic flux through the elementary cell of the ion lattice is small in comparison with the quantum unit. Taking into account the value of the physical parameters in the real single crystal normal metal (like gold, for example) we have finally $1t \ll B \ll 10^3 t$ for the temperatures like $T \leq 1K$. We will not discuss here any questions of rigorous foundations of this approach (very standard in the physics literature dedicated to the transport phenomena). The detailed explanations of this method can be found in classical books (see, for example [7, 8, 9, 10]). Let us give here also the references [35, 36] where the mathematically rigorous approach to the semiclassical motion of electron in electromagnetic field and lattice as well as the historical remarks can be found. Indeed, no rigorous theory of the Kinetic equation here was developed yet, so these papers don't make our results about conductivity more rigorous.

We will consider the electron states in crystal parameterized by the energy bands and the quasimomentum $\mathbf{p}$ defined modulo the reciprocal lattice vectors. From the topological point of view we can say that quasimomentum belongs to the three-dimensional torus $\mathbb{T}^3$ (Brillouen zone) rather than to the Euclidean space $\mathbb{R}^3$. The torus $\mathbb{T}^3$ arises as factorization of the space $\mathbb{R}^3$ with respect to the reciprocal lattice. Topologists say that the space $\mathbb{R}^3$ is a covering over 3-torus $\mathbb{T}^3$. The periodic dispersion relation $\epsilon(\mathbf{p})$ of any energy band can be considered as the one-valued continuous function on the torus $\mathbb{T}^3$. The Fermi surface $S_F : \epsilon(\mathbf{p}) = \epsilon_F$ can also be considered as the smooth compact two-dimensional surface without boundary embedded in the three-dimensional torus $\mathbb{T}^3$. In this paper we will often compare these two pictures

in the 3-torus $\mathbb{T}^3$ and in the total Euclidean 3-space of quasimomenta. We will use the equation $\dot{\mathbf{p}} = \mathbf{F}_{ext}$ both in the torus $\mathbb{T}^3$ and in the covering Euclidean 3-space $\mathbb{R}^3$ for the homogeneous force $\mathbf{F}_{ext}$. In particular, we will consider the properties of the trajectories of this system both in these two spaces which will be very convenient for our consideration. We apologize for using topological terminology in some parts of this paper. However we could not avoid it.

Following the standard approach, we consider a system:

$$\dot{\mathbf{p}} = \frac{e}{c} [\nabla \epsilon(\mathbf{p}) \times \mathbf{B}] + e\mathbf{E}$$

for the semiclassical electron in both homogeneous electric and magnetic field. The value of electric field E is going to be infinitesimally small in measuring the conductivity. Therefore only the trajectories of the main dynamical system

$$\dot{\mathbf{p}} = \frac{e}{c} [\nabla \epsilon(\mathbf{p}) \times \mathbf{B}] \tag{1}$$

should be investigated in this approach.

The trajectories of (1) in the Euclidean 3-space are given on every energy level $\epsilon(\mathbf{p}) = const$ by the plane sections orthogonal to magnetic field. So we have the analytic integrability of the system (1) in the 3-space $\mathbb{R}^3$. However, the global structure of the trajectories on the 3-torus can be highly nontrivial after identification the quasimomenta equivalent modulo the reciprocal lattice.

The dynamical system (1) conserves also the volume element d^3p in $\mathbb{T}^3$ and does not change at all the Fermi distribution. So, in the absence of the electric field $\mathbf{E}$ we will have the electron distribution unchanged (up to the quantum corrections). Nevertheless, the response of this system to small perturbations will be completely different from the case $\mathbf{B} = 0$ and depend strongly on the geometry of trajectories of the dynamical system (1).

Let us say that this dependence was first discovered by the school of I.M. Lifshitz (I.M. Lifshitz, M.Ya. Azbel, M.I. Kaganov, V.G. Peschanskii [1, 2, 3, 5, 6, 8]) in 1950's. Thus, in the work [1] the crucial difference in conductivity was found for the contribution of the closed and open periodic electron trajectories in $\mathbf{p}$ -space considered as the total Euclidean 3-space $\mathbb{R}^3$. Namely, it was shown that the first case corresponds to the total decreasing of conductivity in the plane orthogonal to $\mathbf{B}$ for $B \rightarrow \infty$ while the second case

corresponds to the strong anisotropy of conductivity in the plane orthogonal to $\mathbf{B}$ in the same limit: the conductivity vanishes just in one direction in this plane only depending on the mean direction of the open periodic trajectory. In the works [2, 3] the interesting examples of Fermi surfaces and electron trajectories were considered. However the work [3] contains some conceptual mistake: open trajectories were found for the generic family of magnetic fields with different mean directions. This result is wrong. It contradicts to the "Topological Resonance" which is a base of our main results [23, 28]. We will discuss it in the Chapter 2.

The problem of classification of all possible trajectories on the Fermi surfaces was first set by S.P. Novikov ([11]) and then considered in his school (S.P. Novikov, A.V. Zorich, I.A. Dynnikov, S.P. Tsarev, A.Ya. Maltsev [12, 13, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34]).

The full classification of all possible situations in the Geometric Strong Magnetic Field Limit (GSMF-limit) can be given as a result of the topological studies of this important class of dynamical systems on 2-dimensional surfaces. The most important feature of this new picture is the invention of the observable "Topological numbers" in the conductivity which always appear in GSMF-limit in the situation when the conductivity in the plane orthogonal to the generic magnetic field $\mathbf{B}$ reveals the strong anisotropy for $B \rightarrow \infty$ which is stable with respect to the small rotations of the directions of $\mathbf{B}$. These Topological numbers have the form of the triples of integers $(m_1^\alpha, m_2^\alpha, m_3^\alpha)$. They are connected with some integral planes Γ^α in the reciprocal lattice. The corresponding directions of $\mathbf{B}$ for which the given triple $(m_1^\alpha, m_2^\alpha, m_3^\alpha)$ can be observed give always a region Ω_α of non-zero measure (on the unit sphere) among the total set of directions of $\mathbf{B}$. We call the corresponding region Ω_α on the unit sphere the "Stability zone" corresponding to given triple $(m_1^\alpha, m_2^\alpha, m_3^\alpha)$ due to the topological "rigidity" of the triple $(m_1^\alpha, m_2^\alpha, m_3^\alpha)$ within Ω_α .

We claim also that there are only two types of the stable conductivity tensor asymptotic behavior in the GSMF-limit ($B \rightarrow \infty$) for any normal metal with arbitrary topologically complicated Fermi surface. Namely, the ("topologically nontrivial") case of the strongly anisotropic behavior of conductivity in the plane orthogonal to $\mathbf{B}$ corresponding to some triple of Topological numbers and the ("trivial") case of the uniform decreasing of conductivity in any direction orthogonal to $\mathbf{B}$ for $B \rightarrow \infty$. These cases cover the area on the 2-sphere of the full total measure, so the generic directions are either of the first type or of the second type.

All other types of conductivity behavior in GSMF-limit can not be stable with respect to small rotations of $\mathbf{B}$. We don't give in this paper all topological proofs of these facts because of their rather high mathematical complexity and give just the corresponding citations on the mathematical and physical literature ([11, 12, 13, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34]). Let us introduce here two notations for these stable situations which we will use in this paper.

Situation A.(Topologically trivial behavior).

We call it the situation A; it is the case of uniform decreasing of conductivity in the plane orthogonal to $\mathbf{B}$ for $B \rightarrow \infty$.

Situation B. (Topological numbers and Topological Resonance).

This is the case of the strong anisotropy of conductivity in the plane orthogonal to $\mathbf{B}$ with decreasing in just one direction in this plane for $B \rightarrow \infty$. This direction can be described as the intersection of the plane orthogonal to magnetic field with some integral plane (given by two reciprocal lattice vectors). The corresponding integral plane remains unchanged under the small rotations of the magnetic field. Three integers characterizing this plane in the reciprocal lattice are exactly the observable topological numbers. Topological Resonance claiming that the mean directions of all open trajectories coincide for the generic magnetic field is a base of this result. It was extracted by the present authors from the core of the topological works quoted above. As it was already mentioned, the conceptual mistake has been made exactly here in the classical works of the Lifshitz group.

As we just said above all the stable cases in GSMF-limit can be described just by two situations. However, for complicated enough Fermi surfaces also the rather nontrivial "chaotic" behavior of conductivity tensor in GSMF-limit is possible ([27, 28, 32, 33]) for the set of directions of the zero measure. The trajectories of this type were completely unknown in classical literature. They were discovered recently in the topological works ([19, 25, 29]). Chaotic trajectories can be divided into two different classes:

- 1)Weakly chaotic trajectories (the Tsarev type);
- 2)Strongly chaotic trajectories (the Dynnikov type).

Let us say here some words about these two classes.

The trajectories of the first kind can appear only if the direction of $\mathbf{B}$ is "partly rational", i.e. the plane $\Pi(\mathbf{B})$ orthogonal to $\mathbf{B}$ contains one (up to the multiplier) reciprocal lattice vector. The trajectories of the second kind can appear only if the direction of $(\mathbf{B})$ is completely irrational, i.e. $\Pi(\mathbf{B})$ does not contain any reciprocal lattice vectors. In the case when the direction of

$\mathbf{B}$ is purely rational (i.e. $\Pi(\mathbf{B})$ contains two linearly independent reciprocal lattice vectors) the chaotic electron trajectories can not appear.

The corresponding behavior of conductivity in GSMF-limit is also very different for these two classes of chaotic electron trajectories ([27, 28, 32, 33]). Thus in the case of weakly chaotic trajectories the asymptotic expression of conductivity is just slightly different from the Situation B and corresponds also to strongly anisotropic behavior of conductivity in the plane $\Pi(\mathbf{B})$ for $B \rightarrow \infty$ ([32, 33]). Nevertheless, this regime is unstable with respect to the small rotations of $\mathbf{B}$ unlike the regular case where the "Stability zones" can be observed.

The strongly chaotic trajectories, however, demonstrates completely different behavior of $\sigma^{ik}(B)$ in GSMF-limit ([27]). Namely, in this case the conductivity in the plane orthogonal to $\mathbf{B}$ decreases as $B \rightarrow \infty$ (in all directions) with the different from the Situation A analytic dependence on B . Besides that, in this case the part of the Fermi surface is excluded also from the conductivity along the direction of $\mathbf{B}$ for $B \rightarrow \infty$. The last fact leads to the "sharp minimum" in the conductivity along $\mathbf{B}$ for the given direction of $\mathbf{B}$ if the chaotic trajectories of the second kind appear. Usually the conductivity along $\mathbf{B}$ remains finite in this "sharp minima" since only a part of the Fermi surface becomes excluded from the corresponding contribution. However, these minima can be observed (on the unit sphere) using the small rotations of $\mathbf{B}$ due to the instability of all the chaotic trajectories with respect to such rotations.

In Chapter 3 we give more detailed description of the chaotic trajectories with the corresponding consideration of conductivity.

2 Topologically rigid cases and Topological numbers.

Let us define the "Degree of irrationality" of magnetic field with respect to reciprocal lattice.

Let $\{\mathbf{g}_1, \mathbf{g}_2, \mathbf{g}_3\}$ be the basis of the reciprocal lattice Γ^ such that the vectors of Γ^* are given by all possible integer linear combinations of $\{\mathbf{g}_1, \mathbf{g}_2, \mathbf{g}_3\}$. Then:*

1) The direction of $\mathbf{B}$ is rational (or has irrationality 1) if the plane $\Pi(\mathbf{B})$ orthogonal to $\mathbf{B}$ contains two linearly independent reciprocal lattice vectors.

2) *The direction of $\mathbf{B}$ has irrationality 2 if the plane $\Pi(\mathbf{B})$ contains just one (up to multiplier) reciprocal lattice vector.*

3) *The direction of $\mathbf{B}$ has irrationality 3 (or completely irrational) if the plane $\Pi(\mathbf{B})$ does not contain any reciprocal lattice vectors.*

The generic directions of magnetic field are completely irrational. The direction of $\mathbf{B}$ should be "specially chosen" to have irrationality 1 or 2. We are going to consider now the situations stable with respect to the small rotations of $\mathbf{B}$. This means in particular that specific features of such cases should not be connected with any kind of rationality of the direction of $\mathbf{B}$, i.e. they should reveal all their properties for the completely irrational directions of magnetic field.

The electron trajectories are given by the intersections of the periodic Fermi surface with the family of parallel planes orthogonal to the magnetic field. For simplicity we will assume in this Chapter that the direction of $\mathbf{B}$ is completely irrational (for example, no open periodic trajectories can appear in the planes orthogonal to $\mathbf{B}$). Let us postpone the specific (unstable) features of rational directions to the next Chapter.

We call the trajectory non-singular if it is not adjacent to the critical point. The trajectories adjacent to the critical points as well as the critical points themselves we call singular trajectories.

We call the non-singular trajectory compact if it is closed on the plane. We call the non-singular trajectory open if it is unbounded in $\mathbb{R}^2$.

The examples of singular, compact and open non-singular trajectories are shown on the Fig. 1, a-c.

We give now the important definitions concerning the behavior of open trajectories in the planes orthogonal to $\mathbf{B}$. It is this type of trajectories which plays the main role in the GSMF-limit of conductivity.

We call the open trajectory topologically regular (i.e. corresponding to the "topologically integrable" case) if it lies within the straight line strip of the finite width in $\mathbb{R}^2$ and passes through it from $-\infty$ to ∞ (see Fig. 2, a). All other open trajectories we call chaotic (Fig. 2, b).

Let us point out that the topologically regular open trajectories are not periodic at all which would contradict to the irrationality of the direction of $\mathbf{B}$. In fact they are quasiperiodic.¹ The property of topological regularity is connected with the rather nontrivial property of the "Carriers" of such trajectories in the $\mathbf{p}$ -space which we are going to consider below. Let us

¹The ergodic properties of trajectories of this kind were investigated in [14, 15]




Figure 1: The singular, compact and open non-singular trajectories. The signs " + " and " - " show the regions of larger and smaller values of $\epsilon(\mathbf{p})|_{\Pi}$ respectively.




Figure 2: "Topologically regular" (a) and "chaotic" (b) open trajectories in the plane Π orthogonal to $\mathbf{B}$.




Figure 3: The cylinder of compact trajectories bounded by the singular orbits. (The simplest case of just one critical point on the singular trajectory.)

just say now that this property is closely connected with the "Topological numbers" which we are going to introduce.

Let us introduce now the "Carriers of open trajectories" and the "Topological numbers". We follow here the convenient description ([25],[29]) of the Fermi surface with the trajectories on it when the direction of $\mathbf{B}$ is fixed. We will be interested first of all in the open electron trajectories in the $\mathbf{p}$ -space. Let us say that in general just a part of the Fermi surface will be covered by the open electron trajectories. The remaining part will contain compact (or singular) trajectories. Let us remove all parts of the Fermi surface covered by the non-singular compact trajectories. The remaining part

$$S_F / (\text{Compact Nonsingular Trajectories}) = \cup_j S_j$$

is a union of the 2-manifolds S_j with boundaries ∂S_j who are the compact singular trajectories. The generic type in this case is a separatrix orbit with just one critical point like on the Fig. 3.

We call every piece S_j the "Carrier of open trajectories".




Figure 4: The reconstructed constant energy surface with removed compact orbits and with the two-dimensional discs attached to the singular orbits; in the generic case there is just one critical point on every singular orbit.

These pieces of Fermi surface, however, has the holes with boundaries. They are not the "closed manifolds" anymore. To get the closed manifolds let us make the next step:

We fill in the holes by the topological $2D$ discs in the planes orthogonal to $\mathbf{B}$; finally we are coming to the closed surfaces

$$\bar{S}_j = S_j \cup (2Ddiscs)$$

(see Fig. 4).

This procedure gives the periodic surface $\bar{S}_F$ after the reconstruction and we can define the "compactified carriers of open trajectories" both in $\mathbb{R}^3$ and $\mathbb{T}^3$. Thus we have two representations of the reconstructed Fermi surface:

- 1) The compact surface without boundary embedded in the space of quasi-momenta $\mathbb{T}^3$ (consisting of several pieces without boundaries);
- 2) The set of periodic two-dimensional surfaces without boundaries in the covering space $\mathbb{R}^3$.

Let us formulate now our main intermediate result which was established using the theorems extracted from the purely topological investigations (see, for example [12]).

Let us fix the generic direction of $\mathbf{B}$ and consider the set $\{\bar{S}_j\}$ carrying the open electron trajectories. Then the only two situations can be topologically stable with respect to the small rotations of $\mathbf{B}$:

- (A) *The set $\{\bar{S}_j\}$ is empty;*
- (B) *The set $\{\bar{S}_j\}$ in the torus $\mathbb{T}^3$ consists of the even number of surfaces homeomorphic to the two-dimensional tori $\mathbb{T}_j^2$; all of them have the same homology class in $H_2(\mathbb{T}^3)$ up to the sign (sum of these classes is equal to zero). This property was called the "Topological resonance". The corresponding representation of the set $\{\bar{S}_j\}$ in total $\mathbf{p}$ -space $\mathbb{R}^3$ can be described as follows:*

The manifolds $\bar{S}_j$ represent the periodically deformed two-dimensional planes $\Gamma_{(j)\alpha}$ embedded in $\mathbb{R}^3$ with the same common integer-valued mean directions. In other words we have the set of periodically deformed (warped) integral planes in $\mathbb{R}^3$ which are all parallel in average and do not intersect each other. This picture remains unchanged after the small rotation of the magnetic field.

The first situation A corresponds to the absence of the open electron trajectories on the Fermi level. Let us remind also that we call the two-dimensional plane "integral" in $\mathbb{R}^3$ if it is generated by two reciprocal lattice vectors.

The Topological resonance was first pointed out in [23, 28]. It plays the crucial role in the GSMF-limit as we will see below.

The topological stability means in particular that the corresponding picture remains the same after any rotation of $\mathbf{B}$ small enough: the number of connected components as well as the homological classes of corresponding tori $\mathbb{T}_j^2$ are the same for all the directions of $\mathbf{B}$ close enough to the initial one. Let us make now the important physical conclusion from our main statement and consider the corresponding corollaries for the electrical conductivity.

It was also proved ([29]) that the total measure of the directions of $\mathbf{B}$ where different situations can arise is zero on the unit sphere for the generic

Fermi surface S_F .

We will consider these two situations described above as the main basic foundation for the total classification of different regimes in the GSMF-limit for the generic case.

Let us define now the "Topological numbers" observable in the Situation B when we really have regular open trajectories.

We call the "Topological numbers" corresponding to the stable open electron trajectories the triple of integers (m_1, m_2, m_3) representing the integral 2-plane in the 3-space with reciprocal lattice. (Topologically it is a common homology class of the 2-tori $\mathbb{T}_j^2$ in $\mathbb{T}^3$.)

This integers (m_1, m_2, m_3) can be extracted from common directions of periodically deformed two-dimensional planes representing $\{\bar{S}_j\}$ in $\mathbb{R}^3$ with respect to reciprocal lattice Γ^* . Namely, the planes Γ_α can be defined from the equation

$$m_\alpha^1[\mathbf{x}]_1 + m_\alpha^2[\mathbf{x}]_2 + m_\alpha^3[\mathbf{x}]_3 = 0$$

where $[\mathbf{x}]_i$ are the coordinates in the basis $\{\mathbf{g}_1, \mathbf{g}_2, \mathbf{g}_3\}$ of the reciprocal lattice, or equivalently

$$m_\alpha^1(\mathbf{x}, \mathbf{l}_1) + m_\alpha^2(\mathbf{x}, \mathbf{l}_2) + m_\alpha^3(\mathbf{x}, \mathbf{l}_3) = 0$$

where $\{\mathbf{l}_1, \mathbf{l}_2, \mathbf{l}_3\}$ is the basis of the initial lattice in the coordinate space.

We can formulate now the main statement about the stable open trajectories in our approach:

All stable open electron trajectories have the topologically regular form, i.e. lie in the straight strips of the finite width in the planes orthogonal to $\mathbf{B}$ in the $\mathbf{p}$ -space and pass through them. All trajectories of this kind have the same mean directions for the given direction of $\mathbf{B}$: in average they are parallel to each other. The common direction of all these trajectories is given by the intersection of plane $\Pi(\mathbf{B})$ orthogonal to $\mathbf{B}$ with some integral plane Γ_α which is locally stable with respect to the small rotations of $\mathbf{B}$.

The fact that all topologically regular trajectories are parallel to each other expresses here the "Topological Resonance" property. It first appeared in [23, 28]. It seems that nothing like that was known in the classical literature. In the work [3] for example the open electron trajectories with different

mean directions were mistakably demonstrated for some analytic dispersion relations in the whole regions of the unit sphere parameterizing directions of $\mathbf{B}$. We claim however, that this situation is completely impossible for any open region on the sphere. The important property of topologically regular open trajectories lies in the following fact: their contribution to the conductivity does not differ in the main order in the GSMF-limit from the (anisotropic) contribution of open periodic trajectories obtained in the old work [1]. It is very easy to prove this statement taking into account that the motion of electron is linear plus something bounded: one should simply repeat the essential arguments of this old work. The "Topological Resonance" claims more: all trajectories of this kind give the same kind of anisotropy in the same coordinate system. Only this result makes this behavior experimentally observable. Let us present here corresponding expressions for the conductivity in the GSMF-limit for two situations described above.

Case I (Compact trajectories only):

$$\sigma^{ik} \simeq \frac{ne^2\tau}{m^*} \begin{pmatrix} (\omega_B\tau)^{-2} & (\omega_B\tau)^{-1} & (\omega_B\tau)^{-1} \\ (\omega_B\tau)^{-1} & (\omega_B\tau)^{-2} & (\omega_B\tau)^{-1} \\ (\omega_B\tau)^{-1} & (\omega_B\tau)^{-1} & * \end{pmatrix}, \quad \omega_B\tau \rightarrow \infty \quad (2)$$

Case II (Open topologically regular trajectories):

$$\sigma^{ik} \simeq \frac{ne^2\tau}{m^*} \begin{pmatrix} (\omega_B\tau)^{-2} & (\omega_B\tau)^{-1} & (\omega_B\tau)^{-1} \\ (\omega_B\tau)^{-1} & * & * \\ (\omega_B\tau)^{-1} & * & * \end{pmatrix}, \quad \omega_B\tau \rightarrow \infty \quad (3)$$

Here $\simeq$ means "of the same order in $\omega_B\tau$ and $*$ are some constants ~ 1 . We assume here that the z -axis is always directed along the magnetic field $\mathbf{B}$ and the x -axis in the plane $\Pi(\mathbf{B})$ (orthogonal to $\mathbf{B}$) is directed along the common mean direction of the topologically regular trajectories in $\mathbf{p}$ -space in the second case. Let us mention also that the relations (2)-(3) give only the order of magnitude of σ^{ik} .

The anisotropy of the tensor σ^{ik} in the formula (3) gives the experimental possibility of measuring the mean directions of the topologically regular open orbits for rather big values of B . Using the rotations of the direction of

B it is possible also to find the "Stability zone" on the unit sphere and to determine the corresponding "Topological numbers" characterizing this stable situation. We see that there is only one direction $\hat{\eta}$ in the second case where the conductivity vanishes in the limit $B \rightarrow \infty$. According to the formula (3) this direction coincides with the common mean direction of the topologically regular trajectories in the $\mathbf{p}$ -space (i.e. orthogonal to the mean direction of these trajectories in the coordinate $\mathbf{x}$ -space). The direction $\hat{\eta}(\mathbf{B})$ depends on the direction of magnetic field. However, it varies in some integral plane Γ_α which is the same for the given "Stability zone". We can claim that the direction of conductivity decreasing $\hat{\eta} = (\eta_1, \eta_2, \eta_3)$ satisfies to the relation

$$m_\alpha^1(\hat{\eta}, \mathbf{l}_1) + m_\alpha^2(\hat{\eta}, \mathbf{l}_2) + m_\alpha^3(\hat{\eta}, \mathbf{l}_3) = 0$$

for all the points of stability zone Ω_α which makes possible the experimental observation of the numbers $(m_\alpha^1, m_\alpha^2, m_\alpha^3)$.

3 The chaotic cases and their contribution to the GSMF-limit.

Let us say now some words about the chaotic trajectories which can arise in the special cases for rather complicated Fermi surfaces. You have to remember that they can appear only for the zero measure set of directions of the magnetic field. We think that for the generic Fermi surfaces the fractal (or Hausdorf) dimension of this set is strictly less than 1 (it was certainly proved by Dynnikov that it is no more than 1 for the generic Fermi surfaces, but it can be more for the nongeneric ones—see numerical studies in the works [29, 31]. Anyway, there is no proof of this until now.

We will first mention Tsarev's example of weakly chaotic trajectory having an asymptotic direction in $\mathbb{R}^3$ ([19]). We will not describe here the details of corresponding Fermi surface (see [33]) and just say the trajectory of this kind can not be imbedded in any straight strip of finite width in $\mathbf{p}$ -space. However this trajectory has always asymptotic direction in the plane orthogonal to $\mathbf{B}$. The motion is linear plus smaller (but unbounded) terms. We can always choose the coordinate system such that the average values of the group velocities satisfy to the following condition:

$$\langle v_{gr}^x \rangle = 0 \quad , \quad \langle v_{gr}^y \rangle \neq 0 \quad , \quad \langle v_{gr}^z \rangle \neq 0$$

Here again the z -axis coincides with the direction of $\mathbf{B}$ and the x -axis is directed along the asymptotic direction of the chaotic trajectory in $\mathbf{p}$ -space.

The corresponding behavior of conductivity in GSMF-limit does not coincide completely with the formula (3), however the following formulae for the behavior of $\sigma^{ik}(B)$ can be proved:

$$\sigma^{ik}(\mathbf{B}) \simeq \frac{ne^2\tau}{m^*} \begin{pmatrix} o(1) & o(1) & o(1) \\ o(1) & * & * \\ o(1) & * & * \end{pmatrix} \quad , \quad \omega_B\tau \rightarrow \infty \quad (4)$$

which replaces the formula (3) for the case of weakly chaotic trajectories. Let us omit here all details of the weakly chaotic trajectories and just point out that the asymptotic direction of the weakly chaotic trajectory can be also observed experimentally due to the same reasons as in the case of topologically regular trajectories. However, unlike the topologically regular case, the weakly chaotic trajectories are unstable with respect to generic small rotations of $\mathbf{B}$. They correspond to some very small sets on the unit sphere. At last we say that the trajectories of this kind can appear only for the direction of $\mathbf{B}$ of irrationality 2, i.e. the plane $\Pi(\mathbf{B})$ should contain one reciprocal lattice vector in this situation.

Let us say now some words about more general strongly chaotic trajectories which do not have any asymptotic direction in $\mathbb{R}^3$. We will not describe here the corresponding construction (see [25]) and just give the main features of such trajectories.

First of all, these trajectories can arise only in the case of magnetic field of irrationality 3 and the corresponding carriers have then the genus ≥ 3 . This kind of trajectories are completely unstable with respect to the small rotations of $\mathbf{B}$ and can be observed just for special fixed directions of $\mathbf{B}$ in the case of rather complicated Fermi surfaces. The approximate form of some trajectories of this kind is shown at Fig. 2, b. Moreover, if the genus of Fermi surface is not very high (< 6) it can always be stated that the corresponding carrier of open strongly chaotic trajectories is invariant under the involution $\mathbf{p} \rightarrow -\mathbf{p}$ (after the appropriate choice of the initial point in $\mathbb{T}^3$). The ergodic theorem applied to the open trajectories on the carrier gives then immediately the relations:

$$\langle v_{gr}^x \rangle = 0 \quad , \quad \langle v_{gr}^y \rangle = 0 \quad , \quad \langle v_{gr}^z \rangle = 0$$

for all three components of the group velocity on any of such trajectories. This important fact leads to the rather non-trivial behavior of corresponding contribution to the conductivity for $B \rightarrow \infty$. Namely we can show that all the components of the corresponding contribution to $\sigma^{ik}(\mathbf{B})$ become actually zero in the limit $B \rightarrow \infty$ ([27]). We can write then for this contribution:

$$\sigma^{ik}(\mathbf{B}) \simeq \frac{ne^2\tau}{m^*} \begin{pmatrix} o(1) & o(1) & o(1) \\ o(1) & o(1) & o(1) \\ o(1) & o(1) & o(1) \end{pmatrix} \quad (5)$$

for $B \rightarrow \infty$.²

We see that the strongly chaotic trajectories ("Dyannikov type") do not give any contribution even for conductivity along the magnetic field $\mathbf{B}$ for rather big values of B . In the work [27] also the special "scaling" asymptotic behavior of $\sigma^{ik}(\mathbf{B})$ were suggested. Let us note, however, that the full conductivity tensor include also the contribution of compact (closed) electron trajectories having the form (2) which presents in general as the additional contribution in all the cases described above. We can so claim that the strongly chaotic behavior does not remove completely the conductivity along the magnetic field $\mathbf{B}$ because of the contribution of compact trajectories. However, the sharp local minimum in this conductivity can still be observed in this case since a part of the Fermi surface will be effectively excluded from the conductivity in this situation.

It can be proved (see [29]) that for the generic Fermi surfaces the measure of directions of magnetic field $\mathbf{B}$ where the strongly chaotic behavior can be found on the Fermi surface is equal to zero. However, the total set on the unit sphere corresponding to the strongly chaotic trajectories of this kind can be rather complicated set with the non-trivial Hausdorff dimension. We expect that the Hausdorff dimension of this set is strictly less than 1 for the generic Fermi surfaces. For the nongeneric cases it might be even more than 1.

At last let us say that we expect that either the small stability zones or the strongly chaotic trajectories in fact were observed in the experimental

²Actually the component $\sigma^{zz}(\mathbf{B})$ contains the non-vanishing term of order of T^2/ϵ_F^2 for $B \rightarrow \infty$ for non-zero temperatures ([27]). However, this parameter is very small for the normal metals and we don't take it here in the account.

data represented in [4] (see [27, 28]). However these data are not detailed enough (for example the conductivity along magnetic field was not measured in this experiments).

Let us describe now the total picture for the angle diagram of conductivity in normal metal in the case of geometric strong magnetic field limit ([32, 33]. Namely, we can observe the following objects on the unit sphere parameterizing the directions of $\mathbf{B}$:

1) The "stability zones" Ω_α corresponding to topologically regular open trajectories and parameterized by some integral planes Γ_α in the reciprocal lattice ("Topological Numbers"). All the "stability zones" have the piecewise smooth boundaries on S^2 .

The corresponding behavior of conductivity is described by the formula (3) and reveals the strong anisotropy in the planes orthogonal to the magnetic field. For rather complicated Fermi surfaces we can observe also the "sub-boundaries" of the stability zones where the coefficients in (3) can have the sharp "jump" but do not change the "Topological Numbers" characterizing the "Stability zone" Ω_α .

2) The net of the one-dimensional curves containing directions of irrationality ≤ 2 where the additional periodic open trajectories in $\mathbf{p}$ -space can appear. The corresponding parts of the net are always the parts of the big (passing through the center of S^2) circles orthogonal to some reciprocal lattice vector. The asymptotic behavior of conductivity is given again by the formula (3) but unstable with respect to the small rotations of $\mathbf{B}$ going out from the corresponding curves.

3) The "Special rational directions".

We call the special rational direction the direction of $\mathbf{B}$ orthogonal to the integral plane Γ_α corresponding to some stability zone Ω_α in case when this direction belongs to the same stability zone on the unit sphere. We don't discuss here all the specific features which can appear in this situation and just say that some specialties can arise here. Let us give here the reference on the papers [32, 33] where all corresponding possibilities are discussed.

4) The weakly chaotic open orbits ($\mathbf{B}$ of irrationality 2).

We can have points on the unit sphere where the open orbits are weakly chaotic. All open trajectories still have the asymptotic direction in this case

and the conductivity reveals the strong anisotropy in the plane orthogonal to $\mathbf{B}$ as $B \rightarrow \infty$. The B dependence, however is slightly different from the formula (3) in this case.

5) The strongly chaotic open orbits ($\mathbf{B}$ of irrationality 3).

For some points on S^2 we can have the strongly chaotic open orbits on the Fermi surface. At these points the local minimum of conductivity along the magnetic field is expected. The conductivity along $\mathbf{B}$ however remains finite as $B \rightarrow \infty$ in general situation because of the contribution of compact trajectories.

6) At last we can have the open regions on the unit sphere where only the compact trajectories on the Fermi level are present (Situation A). The asymptotic behavior of conductivity tensor is given then by the formula (2).

At the Fig. 5 we show the schematic picture of the regimes described above for different directions of magnetic field $\mathbf{B}$.

Let now point out some new features connected with the "magnetic breakdown" (self-intersecting Fermi surfaces) which can be observed for rather strong magnetic fields. Up to this point it has been assumed throughout that different parts of the Fermi surface do not intersect with each other. However, it is possible for some special lattices that the different components of the Fermi surface (parts corresponding to different conductivity bands) come very close to each other and may have an effective "reconstruction" as a result of the "magnetic breakdown" in strong magnetic field limit. In this case we can have the situation of the electron motion on the self-intersecting Fermi surface such that the intersections with other pieces do not affect at all the motion on one component. (The physical conditions for the corresponding values of B can be found in [8]). In this case the picture described above should be considered independently for all the non-selfintersecting pieces of Fermi surface and we can have simultaneously several independent angle diagrams of this form on the unit sphere.

References

- [1] I.M.Lifshitz, M.Ya.Azbel, M.I.Kaganov. *Sov. Phys. JETP* **4**, 41 (1957).




Figure 5: The schematic representation of possible regimes for the different directions of the magnetic field $\mathbf{B}$ on the unit sphere.

- [2] I.M.Lifshitz, V.G.Peschansky. *Sov. Phys. JETP* **8**, 875 (1959).
- [3] I.M.Lifshitz, V.G.Peschansky. *Sov. Phys. JETP* **11**, 137 (1960).
- [4] Yu.P.Gaidukov. *Sov. Phys. JETP* **10**, 913 (1960).
- [5] I.M.Lifshitz, M.I.Kaganov. *Sov. Phys. Usp.* **2**, 831 (1960).
- [6] I.M.Lifshitz, M.I.Kaganov. *Sov. Phys. Usp.* **5**, 411 (1962).
- [7] C. Kittel. Quantum Theory of Solids. John Wiley & Sons, Inc. NewYork - London, 1963.
- [8] I.M.Lifshitz, M.Ya.Azbel, M.I.Kaganov. Electron Theory of Metals. Moscow, Nauka, 1971. Translated: New York: Consultants Bureau, 1973.
- [9] J.M. Ziman. Principles of the Theory of Solids. Cambridge, At the University Press, 1972.
- [10] A.A.Abrikosov. Fundamentals of the Theory of Metals. "Nauka", Moscow (1987). Translated: Amsterdam: North-Holland, 1998.
- [11] S.P.Novikov. *Russian Math. Surveys* **37**, 1 (1982).
- [12] A.V.Zorich. *Russian Math. Surveys* **39**, 287 (1984).
- [13] S.P.Novikov. Proc. Steklov Inst. Math. 1 (1986).
- [14] V.I.Arnold. *Functional analysis and its applications* bf 25:2 (1991).
- [15] Ya.G.Sinai, K.M.Khanin. *Functional analysis and its applications* bf 26:3 (1992).
- [16] I.A.Dynnikov. *Russian Math. Surveys* **57**, 172 (1992).
- [17] I.A.Dynnikov. *Russian Math. Surveys* **58** (1993).
- [18] I.A.Dynnikov. "A proof of Novikov's conjecture on semiclassical motion of electron." *Math. Notes* **53**:5, 495 (1993).
- [19] S.P.Tsarev. Private communication. (1992-93).

- [20] S.P.Novikov. "Quasiperiodic structures in topology". Proc. Conference "Topological Methods in Mathematics", dedicated to the 60th birthday of J.Milnor, June 15-22, S.U.N.Y. Stony Brook, 1991. Publish of Perish, Houston, TX, pp. 223-233 (1993).
- [21] A.V.Zorich. Proc. "Geometric Study of Foliations" (Tokyo, November 1993)/ ed. T.Mizutani et al. Singapore: World Scientific, 479-498 (1994).
- [22] S.P.Novikov. Proc. Conf. of Geometry, December 15-26, 1993, Tel Aviv University (1995).
- [23] S.P.Novikov, A.Ya.Maltsev. *ZhETP Lett.* **63**, 855 (1996).
- [24] I.A.Dynnikov. "Surfaces in 3-Torus: Geometry of plane sections." Proc.of ECM2, BuDA, 1996.
- [25] I.A.Dynnikov. "Semiclassical motion of the electron. A proof of the Novikov conjecture in general position and counterexamples." American Mathematical Society Translations, Series 2, Vol. 179, Advances in the Mathematical Sciences. Solitons, Geometry, and Topology: On the Crossroad. Editors: V.M.Buchstaber, S.P.Novikov. (1997)
- [26] I.A.Dynnikov, A.Ya.Maltsev. *ZhETP* **85**, 205 (1997).
- [27] A.Ya. Maltsev. *ZhETP* **85**, 934 (1997).
- [28] S.P.Novikov, A.Ya.Maltsev. *Physics-Uspekhi* **41**(3), 231 (1998).
- [29] I.A.Dynnikov. *Russian Math. Surveys* **54**, 21 (1999).
- [30] S.P.Novikov. *Russian Math. Surveys* **54**:3, 1031 (1999).
- [31] R.D.Leo. SIAM Journal on Applied Dynamical Systems, **2**:4, 517-545 (2003).
- [32] A.Ya.Maltsev, S.P.Novikov, ArXiv: math-ph/0301033, Bulletin of Braz. Math. Society, New Series **34** (1), 171-210 (2003).
- [33] A.Ya.Maltsev, S.P.Novikov.// ArXiv: cond-mat/0304471
- [34] A.Ya. Maltsev, Arxiv: cond-mat/0302014
- [35] G. Panati, H. Spohn, S. Teufel. Phys. Rev. Lett. **88**, 250405 (2002).

[36] G. Panati, H. Spohn, S. Teufel. //arXiv: math-ph/0212041